



The invasive ctenophore *Mnemiopsis leidyi* in northern European waters and its potential impact on fisheries

Jaspers, Cornelia

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Jaspers, C. (2012). *The invasive ctenophore Mnemiopsis leidyi in northern European waters and its potential impact on fisheries*. Paper presented at PICES-2012, Hiroshima, Japan. <http://arctic.au.dk/news-and-events/events/show/artikel/arc-seminar-1/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The invasive ctenophore *Mnemiopsis leidyi* in northern European waters and its potential impact on fisheries



Cornelia Jaspers

DTU Aqua, Technical University of Denmark

S7: Jellyfish in marine ecosystems and their interactions with fish and fisheries
PICES 2012, Hiroshima, Japan 18. Oct. 2012

Mnemiopsis leidyi

- Ctenophore (comb jelly)
- Native: East coast of America
- High feeding rates




Invasive comb jelly *Mnemiopsis leidyi*



Mnemiopsis leidyi in the Black Sea



Fish and *M. leidyi* in the Black Sea




Bilio and Nierman 2004 (MEPS)

Over-exploitation of pelagic fish species e.g. anchovy
Severe eutrophication
Complex interactions

M. leidyi sighted in N. Europe in 2005




Mnemiopsis leidyi in Northern Europe





Bornholm Basin / central Baltic

Spatial and temporal overlap with fish eggs



Cod eggs

Mnemiopsis leidyi

- Temperature ($^{\circ}C$, red line)
- Salinity (psu, blue line)
- Oxygen ($ml\ l^{-1}$, green line)




Objective I

Understand feeding interactions between
M. leidyi and Baltic cod eggs & larvae

Due to high temporal and spatial overlap a large negative impact of
M. leidyi on cod is feared.




Functional response cod egg & larvae



Ingestion rate increases up to 8 cod larvae L^{-1} ;
slope for cod eggs is not sign. different from zero


Food selection

Cod egg vs. *Artemia salina*




Food selection

Cod egg vs. *Artemia salina*




Direct effect:
Results

3D video observations



Mnemiopsis (not) feeding on cod eggs



Mnemiopsis feeding on cod larvae


"Normal" capture response
with lobe closure

Mnemiopsis 35mm oral aboral length

Direct effect: Results



Baltic cod (*Gadus morhua callaris*)



Day 1



Day 2



Day 3



Day 4



Day 5



Day 6



Day 7



Day 8



Day 11



Day 2 post hatching



Day 4 post hatching




Day 6 post hatching




Day 9 post hatching

Magnification: Eggs 25x Larvae 12x


Volume-specific clearance as a function of prey age



Volume-specific clearance as a function of prey age




Volume-specific clearance as a function of prey age




Distribution: Results

Adult *M. leidy* abundances




Distribution: Results

Adult *M. leidy* abundances



Distribution: Results

Adult *M. leidy* abundances



Summary



- Very low feeding rates at 7°C
- Passive negative selection of cod eggs
- Low abundances in the southern & central Baltic
- Applying clearance rates to field abundances:
0.13% cod larvae and 0.05% cod eggs d⁻¹

⇒ No direct predation threat to Baltic cod recruits

Limnol. Oceanogr., 56(2), 2011, 431–439
© 2011, by the American Society of Limnology and Oceanography, Inc.
doi:10.4319/limn.2011.56.2.0431

The invasive ctenophore *Mnemiopsis leidyi* poses no direct threat to Baltic cod eggs and larvae

Cornelia Jaspers,^{a,*} Josefin Titelman,^b Lars Johan Hansson,^c Matilda Haraldsson,^d and Christine Røllike Ditlefsen^d

Objective II




Understand the *in situ* reproduction of *M. leidyi* in the Baltic Sea

Due to the high reproductive capacity, *M. leidyi* could establish a large population in the central Baltic leading to food competition with cod recruits.





Indirect effect: Materials



In situ reproduction rates



Size vs. egg production





Size vs. egg production



Indirect effect: Results


In situ reproduction rates



Zooplankton standing stock 6 times higher in the central Baltic



Jaspers et al. 2011 (PLoS ONE)

Population composition




Indirect effect: Results

Salinity dependent reproduction




Example: High saline area




Limfjorden, Denmark






- Regular "Jelly-Blooms"
- 600-800 *M. leidyi* m⁻³
- Population development
- 11,500 eggs ind⁻¹ d⁻¹
- Active recruitment



Limfjorden, Fisheries



Limfjorden, Fisheries



Mnemiopsis leidyi




- Up to now *M. leidyi* is neither a direct nor an indirect problem for fisheries in the central Baltic
- Potential & documented problem in high saline areas (e.g. Limfjorden, Kattegat)



Conclusion & implications

M. leidyi in European waters

2005 first record in northern Europe (Oliveira 2007)



⇒ *M. leidyi* established throughout European waters





Conclusion & implications

M. leidyi in European waters

Questions arise:
2005 first record in northern Europe (Oliveira 2007)

- 1) Long term impact in high saline areas?
- 2) Might *M. leidyi* acquire a salinity tolerance?
- 3) Climate change effects on population development?



Thomas Kiørboe, DTU Aqua, Denmark

Torkel G. Nielsen

Uffe Thygesen

Fabien Lombard, Villefrance sur Mer, France

Russell R. Hopcroft, University of Alaska, USA

Thorsten Reusch, Geomar, Kiel, Germany

Sören Bolte

Jack Costello, Providence College, RI, USA

Sean Colin, Roger Williams University, RI, USA

BAZOOCA partners:

Peter Tiselius, University of Gothenburg, Sweden

Kajsa Tönnesson

Lene Friis Møller

Matilda Haraldsson

Ulf Båmstedt, Umeå University, Sweden

Lars Johan Hansson, SMHI, Gothenburg, Sweden

Josefin Titelman, University of Oslo, Norway

Aino Hosia, Institute of Marine Research, Norway



Deutscher Akademischer Austausch Dienst
German Academic Exchange Service



Thank you
for your attention!

